

Ordering the affine symmetric group

Alain Lascoux

C.N.R.S., Institut Gaspard Monge, Université de Marne-la-Vallée,
5 Bd Descartes, Champs sur Marne, 77454 Marne La Vallée Cedex 2 FRANCE
Alain.Lascoux@univ-mlv.fr

Abstract. We review several descriptions of the affine symmetric group. We explicit the basis of its Bruhat order.

1 Introduction

For a group defined by generators, one has a notion of *Cayley graph* representing the successive generation of elements of the group by product of generators. This is specially interesting in the case of a Coxeter group $(W, \mathcal{S})$, $\mathcal{S}$ being the set of generators, because one has a length function which filters the elements of W according to the minimum length of the words in the generators of W which expresses them (words of minimum length are called *reduced decompositions*).

The Cayley graph of W naturally provides an ordering (the *weak order*) of W : $w' \leq w$ iff any reduced decomposition of w' is the left factor of at least one reduced decomposition of w , in other words, iff there is a path from w' to w in the Cayley graph of W .

However, if one takes the group algebra of W instead of W , and a reduced decomposition $[i, j, \dots, h] := s_i s_j \dots s_h$ of an element w , then one wants the order to be such that any expression of the type $(1 + s_i)(1 + s_j) \dots (1 + s_h)$ has leading term (with respect to the order) w .

In that case, it is clear that the expansion of $(1 + s_i) \dots (1 + s_h)$ involves only the elements obtained by evaluating in W the subwords of $[i, j, \dots, h]$. One easily checks that the set of such elements is independent of the choice of a reduced decomposition of w (though the set of subwords is not).

Let us write $w' \leq w$ (for the *Bruhat order*) iff there exists a decomposition of w' which is a subword of a reduced decomposition of w . Let us also write $[1, w]$ for the interval $\{w' \mid 1 \leq w' \leq w\}$, 1 being the identity element of W .

The above definition of the Bruhat order [2] is not very efficient, since one has much too many subwords of a reduced decomposition of w compared to the cardinality of $[1, w]$.

Ehresmann[4] found another definition, in the case of the symmetric group, by using all the projections $W \xrightarrow{p_i} W/W_i$ onto all the quotient spaces W/W_i (W_i = maximal parabolic, i.e. group generated by the elements of $\mathcal{S} \setminus s_i$). Ehresmann's definition is

$$w' \leq w \Leftrightarrow p_i(w') \leq p_i(w), \quad i = 1, 2, \dots \quad (1)$$

Now, it makes sense because one has an easy description of the Bruhat order on W/W_i and of the projection. In the case of the symmetric group, W/W_i , as an ordered set, can be identified with the space of all partitions, the diagram of which are included in a fixed rectangle (the order on partitions being inclusion of diagrams).

This description has been generalized by Deodhar[3] to all Coxeter groups.

In [12], M.P. Schützenberger and I gave another description. It uses a notion which is valid for any ordered set X . One defines the basis $\mathcal{B}$ of X to be the following subset:

$$b \in \mathcal{B} \Leftrightarrow \exists x \in X, b \text{ is minimum in the complement of } [1, x] . \quad (2)$$

Given $x \in X$, let furthermore $\mathcal{B}(x)$ denote the set $\{b \in \mathcal{B}, b \leq x\}$. We have shown in [LS] that in the case X is finite, then

$$x \leq y \Leftrightarrow \mathcal{B}(x) \subseteq \mathcal{B}(y) . \quad (3)$$

Now, it happens that in the case of a Coxeter group W the set $\mathcal{B}(w)$ is "small" compared to $[1, w]$ or to the set of subwords of a reduced decomposition of $w \in W$.

The bases of the symmetric group and of the hyperoctahedral group, as well as the map $w \mapsto \mathcal{B}(w)$ are given in [12]. Geck and Kim [8] have determined the bases of all finite Coxeter groups.

The construction of bases passes to the quotient spaces W/W_i (and, as we have said, it is sufficient to treat only these spaces, because of (1)). For example, for the symmetric group $\mathfrak{S}_n$, its quotient $\mathfrak{S}_n/\mathfrak{S}_m \times \mathfrak{S}_r$, $n = m + r$, can be identified with the set of partitions contained in m^r , and its basis consists of all (non void) rectangular partitions $i^j: i \leq m, j \leq r$. The elements of the basis have in fact a geometrical meaning, being in correspondence with determinantal varieties [10].

Now, the notion of a basis can be extended to infinite sets X , when X satisfy some mild hypotheses.

Let us take an affine Coxeter group. Then one still has a length function, which allow to filter the group according to length. Let W^ℓ be the set of elements of length $\leq \ell$. If $\ell(w) \leq \ell$, then the full interval $[1, w]$ is contained in W^ℓ . Let $\mathcal{B}^\ell$ be the basis of W^ℓ .

One has that $\mathcal{B} = \cup \mathcal{B}^\ell$ and one can check that

$$w' \leq w \Leftrightarrow \mathcal{B}(w') \subseteq \mathcal{B}(w) . \quad (4)$$

In the following, we shall describe the basis of the affine symmetric group $\tilde{\mathfrak{S}}_n$ and its Bruhat order.

2 The Affine Symmetric Group

Let n be an integer and $\tilde{\mathfrak{S}}_n$ be the quotient of the infinite symmetric group (generated by $s_i, i \in \mathbb{Z}$ under the relations $s_i = s_{i+n}$), $i \in \mathbb{Z}$. In other words,

$\tilde{\mathfrak{S}}_n$ is generated by $s_0, \dots, s_{n-1}$ satisfying the braid relations (with $s_n := s_0$)

$$s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1}, \quad s_i s_j = s_j s_i, \quad |i - j| \not\equiv 1 \pmod{n}, \quad 0 \leq i, j \leq n - 1. \quad (5)$$

The group $\tilde{\mathfrak{S}}_n$ can be thought as acting on infinite vectors $v = [\dots, v_i, \dots]$, $i \in \mathbb{Z}$: the generator s_i transposes in v components v_{i+kn}, v_{i+1+kn} for all $k \in \mathbb{Z}$ simultaneously. Usually, one takes vectors v such that $v_{i+n} = v_i + n$, $i \in \mathbb{Z}$. Associating to the identity element the “vacuum vector” $\triangleright : \triangleright_i = i$, $\forall i \in \mathbb{Z}$, then the orbit of $\triangleright$ is in bijection with $\tilde{\mathfrak{S}}_n$ and one can code any element $\sigma \in \tilde{\mathfrak{S}}_n$ by the image $v = \triangleright \sigma$ of $\triangleright$ under σ , or equivalently by the *window* $[v_1, \dots, v_n]$.

In other words, elements of $\tilde{\mathfrak{S}}_n$ can be coded by vectors in $\mathbb{Z}^n$, starting from $[1, 2, \dots, n]$, with the action

$$\begin{aligned} [\dots v_i, v_{i+1} \dots] s_i &= [\dots v_i, v_{i+1} \dots], \quad 1 \leq i \leq n - 1 \\ [v_1, \dots, v_n] s_0 &= [v_n - n, \dots, v_1 + n]. \end{aligned} \quad (6)$$

One has in fact many other choices to represent $\tilde{\mathfrak{S}}_n$ by infinite vectors satisfying periodicity conditions. One can also modify the definition of “transposing two adjacent components”.

For example, let $c(\triangleright)$ be the infinite vector with all components equal to 0, and let $c(\sigma)$ be the image of $c(\triangleright)$ under the action

$$[\dots c_j, c_{j+1} \dots] s_i = [\dots c_{j+1} + 1, c_j \dots], \quad 1 \leq i \leq n - 1, \quad c_i \leq c_{i+1}, \quad \forall j \cong i \pmod{n}. \quad (7)$$

This action preserves equalities $c_j = c_{j+n}$, and once more, one can restrict vectors to their components $1, \dots, n$. In that set-up, one has

$$[c_1, \dots, c_n] s_0 = [c_n, \dots, c_1 + 1], \quad c_n \leq c_1. \quad (8)$$

Now, it is easy to check that the mapping $\tilde{\mathfrak{S}}_n \ni \sigma \mapsto [c_1, \dots, c_n] \in \mathbb{N}^n$ is a bijection. It can be easily described without passing through a reduced decomposition of σ (cf. [1]).

By restriction, it induces a bijection from $\mathfrak{S}_n$ onto the set of vectors such that $c_1 \leq n - 1, \dots, c_n \leq 0$. Such a vector is called the *Lehmer code* of the corresponding permutation σ [9] and encodes the “inversions” of σ . This is still true in the infinite case (cf. [1], [5]). Let us call *scalar weight* of $c(\sigma)$ the sum $c_1 + \dots + c_n$. Then the above action shows that the scalar weight increases by 1 if length increases, and thus the scalar weight of $c(\sigma)$ is equal to the length $\ell(\sigma)$. Shi [14] gives another description of the length function, directly from the window of σ .

3 Quotient of $\tilde{\mathfrak{S}}_n$ modulo $\mathfrak{S}_n$

Since all s_i 's play a symmetrical rôle, all quotient spaces of $\tilde{\mathfrak{S}}_n$ modulo a maximal parabolic are isomorphic. Let us choose the finite symmetric group $\mathfrak{S}_n$, generated by $s_1, \dots, s_{n-1}$, as a maximal parabolic.

The unique element of minimum length of a class $\sigma \mathfrak{S}_n$ is the unique element ζ of the class such that

$$\ell(\zeta s_i) < \ell(\zeta) \Leftrightarrow i = 0.$$

Therefore, minimum elements are exactly those elements with (strictly) increasing window $[v_1, \dots, v_n] : v_1 < v_1 < \dots < v_n$.

From (2.3) one deduces that codes of such elements are (weakly) increasing vectors $[c_1, c_2, \dots, c_n]$ such that $c_1 = 0$, that is, are partitions of length $\leq n - 1$.

Now, there is a third coding of $\tilde{\mathfrak{S}}_n/\mathfrak{S}_n$ by infinite vectors u , with periodicity $u_{j+n} = u_j - 1, \forall j \in \mathbb{Z}$. It amounts to take origin vector such that $[u_1, \dots, u_n] = [0, \dots, 0]$, and act by

$$\begin{aligned} [\dots u_i, u_{i+1} \dots] &\xrightarrow{s_i} [\dots u_{i+1}, u_i \dots], 1 \leq i \leq n-1, \\ [u_1, \dots, u_n] &\xrightarrow{s_0} [u_n + 1, \dots, u_1 - 1]. \end{aligned} \tag{9}$$

From a vector u in $\mathbb{Z}^n$ such that $u_1 + \dots + u_n = 0$ (call *weight* such a vector), one can recover an element of $\mathfrak{S}_n \backslash \tilde{\mathfrak{S}}_n$ as follows (the quotient is now on the left): one has to sort increasingly the vector $[1 + nu_1, \dots, n + nu_n]$.

For example, $n = 3, u = [7, -4, -3]$ gives the vector $[22, -10, -6] = [1 + 3 \times 7, 2 - 3 \times 4, 3 - 3 \times 3]$ which is sorted into $[-10, -6, 22]$, and is an element of $\tilde{\mathfrak{S}}_3$ of minimum length (of code $[0, 1, 19]$) in its coset modulo $\mathfrak{S}_3$.

B. Leclerc has shown me that it is more convenient, in the theory of roots of affine Weyl groups, to add an “imaginary root” and use “extended weights” which are vectors in $\mathbb{Z}^{n+1}$: one takes for origin $[0, \dots, 0] \in \mathbb{Z}^{n+1}$ and the action is

$$\begin{aligned} [\dots w_i, w_{i+1} \dots] &\xrightarrow{s_i} [\dots w_{i+1}, w_i \dots], 1 \leq i \leq n-1, \\ [w_1, \dots, w_n, w_{n+1}] &\xrightarrow{s_0} [w_n + 1, \dots, w_1 - 1, w_1 - w_n + w_{n+1} - 1]. \end{aligned} \tag{10}$$

A weight $u \in \mathbb{Z}^n$ already has an unnecessary component, since the sum of components is zero : one can write $u = x_1 \alpha_1 + \dots + x_{n-1} \alpha_{n-1}$, the α_i being the simple roots $[1, -1, 0, \dots, 0], \dots, [0, \dots, 0, 1, -1]$.

However, it happens that going to $\mathbb{Z}^{n+1}$ simplifies some constructions, the extra component having combinatorial interpretations.

4 n -Cores and Bruhat order

None of these three codings furnishes a description of the Bruhat order. Fortunately, Fock spaces and crystal graphs indicate a fourth one ([13], [11], [6]).

Now one will represent elements of $\tilde{\mathfrak{S}}_n/\mathfrak{S}_n$ by partitions which are *n-cores*, that is partitions such that one cannot erase a border strip (also called outer ribbon) of length n from their diagram (cf. [7]).

One numbers diagonals of diagrams with the numbers (which are called *colours*) $0, 1, \dots, n-1$ periodically, the main diagonal being of colour 0.

Now, take the diagram of a an *n-core* such that no corner box is of colour i . Then the action of s_i , $i = 0, \dots, n-1$ on this diagram is to add all possible boxes which are of colour i , in such a way that the resulting object is still the diagram of a partition. If there is at least one corner box of colour i , then the action consists in erasing all peripheal boxes of colour i such that one still gets a diagram of a partition.

It is an easy combinatorics to check that from an *n-core* one still gets an *n-core*, and that the operation is involutive. Now, one does not try to check the braid relations, because one takes the classics again ([7] p.78) and one sees that one can use, instead of an *n-core*, still another object which is called *abacus*.

Let $\diamond$ be a (decreasing) partition extended to infinity on the right by 0's: $\diamond = [\diamond_1, \diamond_2, \dots, \diamond_k, 0, 0, \dots]$. Let $\mathcal{A}$ be the set of numbers $\{\diamond_1 - 0, \diamond_2 - 1, \diamond_3 - 2, \dots\}$ interpreted as balls in an abacus of width n and bi-infinite height, with places numbered consecutively as follows:

$$\begin{array}{cccccc} \vdots & \dots & \vdots & \vdots & & level \\ 1-2n & \dots & -n-1 & -n & -1 & \\ 1-n & \dots & 1 & 0 & 0 & \\ 1 & \dots & n-1 & n & 1 & \\ n+1 & \dots & 2n-1 & 2n & 2 & \\ \vdots & \dots & \vdots & \vdots & \vdots & \end{array}.$$

Putting balls in positions belonging to $\mathcal{A}$ gives by definition the *n-abacus* of the partition $\diamond$.

That $\diamond$ be an *n-core* is equivalent to the fact that balls are vertically packed upwards ([7] p.80).

Reading levels of bottom balls, one gets a vector in $u \in \mathbb{Z}^n$ such that $u_1 + \dots + u_n = 0$, that is, one gets a weight. This construction is, indeed, a bijection between weights in $\mathbb{Z}^n$ and *n-cores* or elements of $\mathfrak{S}_n \backslash \tilde{\mathfrak{S}}_n$ or of $\tilde{\mathfrak{S}}_n/\mathfrak{S}_n$.

For example, the 5-core $[4, 2^3, 1^4]$ gives by subtraction the infinite vector

$$[4, 1, 0, -1, -3, -4, -5, -6, -8, -9, \dots]$$

and the abacus

					<i>level</i>
-14	-13	-12	-11	-10	-2
-9	-8	.	-6	-5	-1
-4	-3	.	-1	0	0
1	.	.	4	.	1

the weight $[1, 0, -2, 1, 0]$, the code $[0, 1, 2, 2, 4]$ and the element $[-7, 2, 5, 6, 9]$ of $\tilde{\mathfrak{S}}_5/\mathfrak{S}_5$.

For each of the different objects that have been given, the description of the action of the simple generators s_i allows to prove that the above constructions are indeed bijections and that the elementary operations represent the generators of $\tilde{\mathfrak{S}}_n$.

Now each of the objects that we have displayed will provide informations about the corresponding element of the affine group, information which would sometimes not have been easy to get in another description, though all bijections here are elementary. For example, it is amusing to look at what becomes the addition of weights (plain addition of vectors) in terms of n -cores, or of codes.

Here is some example computed with ACE [15], where we give a chain of elements of $\tilde{\mathfrak{S}}_4/\mathfrak{S}_4$, with their code, the corresponding extended weight in $\mathbb{Z}^5$ and the 4-core with its colours. The reader should beware that simple transpositions act from the left on codes and on group elements, and from the right on weights and cores.

<i>Code</i>	$[0, 1, 2, 7]$		$[0, 2, 2, 7]$	
<i>Group element</i>	$[-6, 1, 4, 11]$		$[-7, 2, 4, 11]$	
<i>Ext. weight</i>	$[0, -2, 2, 0, -4]$		$[-2, 0, 2, 0, -4]$	
<i>Core</i>	$[7, 4, 2, 2, 1, 1, 1]$		$[7, 4, 2, 2, 2, 1, 1, 1]$	
			1	
	2	$\xrightarrow{s_1}$	2	$\xrightarrow{s_3}$
	3		3	
<i>Diagram</i>	0		0 1	
	1 2		1 2	
	2 3		2 3	
	3 0 1 2		3 0 1 2	
	0 1 2 3 0 1 2		0 1 2 3 0 1 2	

[0, 2, 2, 8]	[0, 2, 2, 9]	[0, 2, 3, 9]
[-7, 2, 3, 12]	[-8, 2, 3, 13]	[-9, 2, 4, 13]
[-2, 0, 0, 2, -4]	[3, 0, 0, -3, -9]	[3, 0, -3, 0, -9]
[8, 5, 2, 2, 2, 1, 1, 1]	[9, 6, 3, 2, 2, 2, 1, 1, 1]	[9, 6, 3, 3, 2, 2, 2, 1, 1, 1]
	3	
	0	
1	1	1
2	2	2 3
3	3 0	3 0
0 1	0 1	0 1
1 2	1 2	1 2 3
2 3	2 3 0	2 3 0
3 0 1 2 3	3 0 1 2 3 0	3 0 1 2 3 0
0 1 2 3 0 1 2 3	0 1 2 3 0 1 2 3 0	0 1 2 3 0 1 2 3 0

For what concerns the Bruhat order, we shall see that we need n -cores. Codes, which are also (increasing) partitions for minimum elements in their coset modulo $\mathfrak{S}_n$, would not give (by inclusion) the proper order, but only a sub-order. Björner and Brenti ([1], th 6.3 and th.6.5) obtained a criterium describing the Bruhat order in terms of monotonous functions; it is equivalent to the following proposition.

Let us denote by $\mathcal{C}$ the morphism from $\tilde{\mathfrak{S}}_n/\mathfrak{S}_n$ (identified with the set of elements of $\tilde{\mathfrak{S}}_n$ which are of minimum length in their coset) onto n -cores.

Proposition 1. *Let $\sigma, \omega \in \tilde{\mathfrak{S}}_n/\mathfrak{S}_n$. Then $\sigma \leq \omega$ iff the diagram of $\mathcal{C}(\sigma)$ is contained in the diagram of $\mathcal{C}(\omega)$.*

Proof. Take any corner of the diagram of $\mathcal{C}(\omega)$ and let i be its colour. Then $s_i \omega < \omega$ and one knows that ([2], [3])

$$\sigma \leq \omega \Leftrightarrow \min(s_i \sigma, \sigma) \leq s_i \omega$$

Now, we have to compare two n -cores with less boxes, and as in the finite case, we are finally reduced to know how to compare two elements of length differing by 1. $\square$

5 Basis of the Bruhat order

When one needs to store huge sets of elements of the affine group, together with the information about the Bruhat order, then it is not efficient to also store the associated partitions (for an element in $\tilde{\mathfrak{S}}_n$, we need n partitions, because we have n different projections $\tilde{\mathfrak{S}}_n \mapsto \tilde{\mathfrak{S}}_n/\mathfrak{S}_n$). It is well illustrated by Geck and Kim [8], in the finite case, that one has instead to use the basis of the order, as defined above. In this section we shall determine the basis of $\tilde{\mathfrak{S}}_n/\mathfrak{S}_n$.

Let us first introduce some special n -cores.

Given two strictly positive numbers p, q such that $p + q = n$, a (p, q) -staircase is a partition E such that the stairs of its diagram are all of height p and width q , except for the top stair which is of width $\leq q$ and the bottom stair which is of height $\leq p$.

In other words, $E = [E_1, E_2, \dots, E_k, 0, \dots]$, the non zero differences $E_i - E_{i+1}$ are all equal to q , except possibly for the last one $E_k - 0$ which is such that $1 \leq E_k \leq q$, and the partition conjugate to E (obtained by transposing the axes) satisfies the same conditions with p instead of q .

Let us notice that all the corners of a staircase have the same colour, and that, given (p, q) and a point in the Cartesian quadrant $\mathbb{N} \times \mathbb{N}$, there exists one and only one (p, q) -staircase having this point as a corner.

Let $\mathcal{D}$ be the diagram of an n -core, and $\triangle$ one of its corners. ∇ the maximal box of $\mathcal{D}$ on the first diagonal of the same colour above (resp. below) the diagonal containing $\triangle$ (if it exists). Define the *left-rectrix* (resp. *right-rectrix*) of $\mathcal{D}$ of *pivot* $\triangle$ to be the staircase having $\triangle$ and ∇ as corners. n -Cores which are too small to have two diagonals of the same colour are in fact staircases. Let us say in that case that they have only one rectrix which coincide with themselves.

For the 4-core $[5, 4, 3, 2, 1, 1, 1]$, here is an example of a left-rectrix (figured by $\heartsuit$'s):

$$\begin{array}{ccccc}
 2 & & 2 & & 2 \\
 3 & & 3 & & 3 \\
 0 & & \nabla & & \heartsuit \\
 1\ 2 & \mapsto & 1\ 2 & \mapsto & \heartsuit\ 2 \\
 2\ 3\ 0 & & 2\ 3\ \triangle & & \heartsuit\ \heartsuit\ \heartsuit \\
 3\ 0\ 1\ 2 & & 3\ 0\ 1\ 2 & & \heartsuit\ \heartsuit\ \heartsuit\ 2 \\
 0\ 1\ 2\ 3\ 0 & & 0\ 1\ 2\ 3\ 0 & & \heartsuit\ \heartsuit\ \heartsuit\ \heartsuit\ \heartsuit
 \end{array}$$

Recall that in the finite symmetric group case, cosets representatives are diagrams contained in a rectangle; rectrices of a diagram $\mathcal{D}$ are the maximal rectangular partitions contained in it [12]. Describing Bruhat order on a finite symmetric group essentially reduces to decomposing partitions into maximal rectangles.

Lemma 2. *Let $\mathcal{D}$ be the diagram of an n -core. Then $\mathcal{D}$ is the supremum (with respect to inclusion of diagrams) of its rectrices.*

Proof. Since $\mathcal{D}$ is contained in the supremum of any family of diagrams such that each corner of $\mathcal{D}$ is contained into at least one of them, it is sufficient (and immediate) to check that each rectrix of $\mathcal{D}$ is contained entirely in $\mathcal{D}$. $\square$

In order to determine the basis of $\tilde{\mathfrak{S}}_n/\mathfrak{S}_n$, we need some more properties of staircases.


Lemma 3. Let $\mathcal{D}$ and $\mathcal{F}$ be diagrams of n -cores such that $\mathcal{D}$ is not a staircase and $\mathcal{D}$ is not included into $\mathcal{F}$. Then there exists at least one staircase E in $\mathcal{D}$ such that $E \not\subset \mathcal{F}$.

Proof. Take any corner $\clubsuit$ of $\mathcal{D}$ which does not belong to $\mathcal{F}$. Then any staircase with corner $\clubsuit$ and included into $\mathcal{D}$ will do. $\square$


Lemma 4. Let E be (p, q) -staircase, having at least one internal corner $\clubsuit$ (i.e. a corner which is not on the first column, nor on the bottom row). Let ∇ , $\triangle$ be the east and south neighbour boxes of $\clubsuit$, and let E_{∇} be the $(n-1, 1)$ -staircase of corner ∇ and $E_{\triangle}$ be the $(1, n-1)$ -staircase of corner $\triangle$. Let moreover F be the partition which is the supremum of E_{∇} and of $E_{\triangle}$ (it is an n -core).
Then E is minimum in the complementary of the interval $[\emptyset \leq F]$.

Proof. Suppose that an n -core E' is strictly contained in E and not in F . Take a box of E' which does not belong to F and a staircase E'' contained in E' of corner this box. Then E'' is not included in F . Therefore it is sufficient to prove that each staircase E'' strictly contained in E is also contained in F . If $E'' \subset E$ contains a box which is an internal corner of E , then this box is a corner of E'' and $E'' = E$. Therefore E'' avoids all internal corners of E , in particular $\clubsuit$. But it is clear that the part of E'' which is right of $\clubsuit$ is contained in $E_{\triangle}$, and that the part which is left of $\clubsuit$ is contained in E_{∇} . This implies that $E'' \subseteq F$. $\square$

For example, take the $(2, 2)$ -staircase $\overset{\circ}{\circ}\overset{\circ}{\circ} := [7, 5, 5, 3, 3, 1, 1]$, choosing the corner pointed by $\clubsuit$, its diagonal being figured by $\spadesuit$'s:



Then any 4-core strictly contained in $\overset{\circ}{\circ}\overset{\circ}{\circ}$ is also contained in the 4-core



We have left aside the cores having no internal corner, that is *hook*-partitions. A case-by-case analysis which offers no difficulty gives the following classification.

Lemma 5. *Partitions*

$$[1]; [2], \dots, [n-1]; [1^2], \dots, [1^{n-1}]; [n, 1], [2, 1^{n-1}]$$

belong to the basis, but not the other hooks with $n+1$ boxes: $[3, 1^{n-2}], \dots, [n-1, 1^2]$, though they are (p, q) -staircases.

For example, for $n = 5$, the basis contains the hooks $[1]; [2], [3], [4]; [1^2], [1^3], [1^4]; [5, 1], [2, 1^3]$, but not $[3, 1^3]$ nor $[4, 1^2]$. Indeed, $[3, 1, 1, 1]$ is the supremum of $[3]$ and $[1, 1, 1, 1]$, and $[4, 1, 1]$ is the supremum of $[4]$ and $[1, 1, 1]$.

In summary, one has the following characterization of the basis of $\tilde{\mathfrak{S}}_n/\mathfrak{S}_n$.

Theorem 6. *The basis of $\tilde{\mathfrak{S}}_n/\mathfrak{S}_n$ for the Bruhat order consists in all (p, q) -staircases, $p + q = n$, excepted the hooks $[3, 1^{n-2}], \dots, [n-1, 1, 1]$.*


We have now all the necessary tools to handle the affine symmetric group together with its Bruhat order. In particular, elements will be represented by the decomposition of their associated cores into maximal sub-staircases.

For example, the 5-core $\overset{\circ}{\circ} = [6, 6, 5, 4, 3^4, 2^4, 1^4]$, corresponding to the group element $[-15, 2, 4, 11, 13]$, with code $[0, 3, 3, 7, 8]$ and length 21, is decomposed into the following three staircases


$$\begin{array}{ccccccc} 0 & & 0 & & & & \\ 1 & & 1 & & & & \\ 2 & & 2 & & & & \\ 3 & & 3 & & & & \\ 40 & & 40 & & & & \\ 01 & & 01 & & & & \\ 12 & & 12 & & & & \\ 23 & & 23 & & & & \\ 340 & = & 340 & \cup & & \cup & \\ 401 & & 401 & & & & \\ 012 & & 012 & & 012 & & \\ 123 & & 123 & & 123 & & \\ 2340 & & 2340 & & 234 & & 234 \\ 34012 & & 3401 & & 34012 & & 340 \\ 401234 & & 4012 & & 40123 & & 401234 \\ 012340 & & 0123 & & 01234 & & 012340 \end{array}$$

For a 5-core to be bigger than $\overset{\circ}{\circ}$ with respect to the Bruhat order, it is necessary and sufficient to be bigger than each of the above three staircases.


As in the case of a finite symmetric group, one has to reduce the numbers of projections $\tilde{\mathfrak{S}}_n \mapsto \tilde{\mathfrak{S}}_n/\mathfrak{S}_n$ or $\tilde{\mathfrak{S}}_n \mapsto \mathfrak{S}_n \setminus \tilde{\mathfrak{S}}_n/\mathfrak{S}_n$ to get an efficient coding [12]. We shall not go into these technicalities here. Let us just mention that in the case of a finite symmetric group, we have used such a “decomposition” of a permutation into rectangular partitions to compute Kazhdan-Lusztig polynomials [10].




Cosets representatives for $n=3$



Codes



Extended Weights



3-Cores

References

1. A. Björner, F. Brenti, *Affine permutations of type A*, Electronic J. Comb. **3**# R18 (1996).
2. N. Bourbaki, *Groupes et Algèbres de Lie*, Fasc 34, Hermann, Paris (1968).
3. V.V. Deodhar, *Some characterizations of Bruhat ordering on a Coxeter group*, Invent. Math. **39** (1977) 187-198.
4. C. Ehresmann, *Sur la topologie de certains espaces homogènes*, Ann. Math. **35** (1934) 396-443.
5. H. Eriksson, *Computational and combinatorial aspects of Coxeter groups*, thesis, KTH, Stockholm (1994).
6. O. Foda, B. Leclerc, M. Okado, J-Y Thibon, T. Welsh, *Branching functions of $A_{n-1}^{(1)}$ and Jantzen-Seitz problem for Ariki-Koike algebras*, Adv. in Maths **141** (1999) 322-365.
7. James, A. Kerber, *The Representation Theory of the Symmetric Group*, Encyclopedia of Mathematics, Addison-Wesley, Reading MA, (1991).
8. M. Geck, S. Kim, *Bases for the Bruhat-Chevalley order on all finite Coxeter groups*, J. of Algebra **197** (1997) 278-310.
9. A. Kerber, *Algebraic Combinatorics via finite group actions*, Wissenschaftsverlag, Mannheim (1991).
10. A. Lascoux, , *Ordonner le groupe symétrique: pourquoi utiliser l'algèbre de Iwahori-Hecke ?*, Congrès International Berlin 1998, Documenta Mathematica, extra volume ICM 1998, III (1998) 355-364.
11. A. Lascoux, B. Leclerc et J.Y. Thibon, *Crystal graphs and q -analogues of weight multiplicities for root systems of type A_n* , Letters in Math. Physics, **35** (1995) 359-374.
12. A. Lascoux, M.P. Schützenberger, *Treillis et bases des groupes de Coxeter*, Electronic J. of Comb. **3** # R27 (1996).
13. K.C. Misra, T. Miwa, *Crystal base of the basic representation of $U_q(\widehat{sl}_n)$* , Commun. Math. Phys., **134**, 1990, p.79-88.
14. Jian-Yi Shi, *The Kazhdan-Lusztig cells in certain affine Weyl Groups*, Springer L.N. **1179** (1986).
15. S. Veigneau, *ACE, an algebraic environment for the computer algebra system MAPLE*, <http://phalanstere.univ-mlv.fr/~ace> (1998).